



**UNIVERSIDADE FEDERAL DO OESTE DO PARÁ**  
**INSTITUTO DE CIÊNCIAS E TECNOLOGIA DAS ÁGUAS**  
**PROGRAMA DE PÓS-GRADUAÇÃO EM BIODIVERSIDADE**

**MARIA KATIANE SOUSA COSTA**

**RESPOSTAS DAS ASSEMBLEIAS DE BESOUROS ESCARABEÍNEOS ÀS  
MUDANÇAS AMBIENTAIS E PRESENÇA DE SILVICULTURA NA AMAZÔNIA  
ORIENTAL**

**SANTARÉM – PA**

**2021**



**UNIVERSIDADE FEDERAL DO OESTE DO PARÁ**  
**INSTITUTO DE CIÊNCIAS E TECNOLOGIA DAS ÁGUAS**  
**PROGRAMA DE PÓS-GRADUAÇÃO EM BIODIVERSIDADE**

**MARIA KATIANE SOUSA COSTA**

**RESPOSTAS DAS ASSEMBLEIAS DE BESOUROS ESCARABEÍNEOS ÀS  
MUDANÇAS AMBIENTAIS E PRESENÇA DE SILVICULTURA NA AMAZÔNIA  
ORIENTAL**

Dissertação apresentada ao Programa de Pós-Graduação em Biodiversidade da Universidade Federal do Oeste do Pará, como requisito para obtenção de grau de Mestre em Biodiversidade.

Orientador: Prof. Dr. Rodrigo Ferreira Fadini  
**Universidade Federal do Oeste do Pará**

Coorientador: Prof. Dr. Filipe Machado França  
**University of Bristol, School of Biological Sciences**

**SANTARÉM – PA**

**2021**

**Dados Internacionais de Catalogação-na-Publicação (CIP)**  
**Sistema Integrado de Bibliotecas – SIBI/UFOPA**

---

C837r Costa, Maria Katiane Sousa

Respostas das assembleias de Besouros Escarabeínos às mudanças ambientais e  
presença de silvicultura na Amazônia Oriental./ Maria Katiane Sousa Costa. – Santa-  
ré, 2021.

48 p. : il.  
Inclui bibliografias.

Orientador: Rodrigo Ferreira Fadini

Coorientador: Filipe Machado França

Dissertação (Mestrado) – Universidade Federal do Oeste do Pará, Pró-reitoria de Pes-  
quisa, Pós Graduação e Inovação Tecnológica, Programa de Pós-Graduação em Biodiver-  
sidade.

1. Brasil. 2. Cobertura do dossel. 3. Efeito de borda. I. Fadini, Rodrigo Ferreira, *ori-ent.* II. França, Filipe Machado, *coorient.* III. Título.

CDD: 23 ed. 632.76098115

---

Bibliotecária - Documentalista: Renata Ferreira – CRB/2 1440



*Universidade Federal do Oeste do Pará*

**PROGRAMA DE PÓS GRADUAÇÃO EM BIODIVERSIDADE**

**ATA Nº 23**

Em acordo com o Regimento do Programa de Pós Graduação em Biodiversidade da Universidade Federal do Oeste do Pará, a dissertação de mestrado é julgada por uma Banca Avaliadora não presencial, constituída por cinco avaliadores, sendo um deles obrigatoriamente externo ao curso, com título de doutor (Artigo 56 do referido regimento). O acadêmico é considerado aprovado quando ao menos três membros avaliadores emitirem pareceres Aprovado. Alternativamente, o discente será dispensado da banca avaliação da dissertação, quando comprovar o aceite ou publicação de pelo menos um artigo resultante da sua dissertação, como primeiro autor, em co-autoria com orientador, ou orientador e coorientador quando o orientador for um docente colaborador, em periódico indexado com percentil mínimo de 75 (setenta e cinco) ou superior referente às métricas mais recentes do maior percentil utilizado pelo Journal Citation Reports (Clarivate) ou pelo Scientific Journal Rankings (Scimago), cabendo ao discente apenas a apresentação pública do trabalho (Artigo 58). O discente que teve sua dissertação aprovada deverá apresentá-la em sessão pública com duração de até 50 (cinquenta) minutos obrigatoriamente até no máximo 15 (quinze) dias após a aprovação, e no prazo máximo de vínculo com o curso, ou seja, 24 (vinte e quatro) meses após o início do primeiro semestre letivo do discente no curso (artigo 64). Assim, aos vinte e sete dias do mês de janeiro do ano de dois mil e vinte e dois, às dezesseis horas, de forma remota através da plataforma GoogleMeet, instalou-se a apresentação de seminário público da dissertação de mestrado da aluna MARIA KATIANE SOUSA COSTA. Deu-se início a abertura dos trabalhos, onde o Professor Dr. Rodrigo Ferreira Fadini, após esclarecer as normativas de tramitação da defesa e seminário público, de imediato solicitou a candidata que iniciasse a apresentação da dissertação, intitulada "RESPOSTAS DAS ASSEMBLEIAS DE BESOUROS ESCARABEÍNEOS ÀS MUDANÇAS AMBIENTAIS E PRESENÇA DE SILVICULTURA NA AMAZÔNIA ORIENTAL". Concluída a exposição, o professor comunicou a discente que a versão final da dissertação deverá ser entregue ao programa, no prazo de 60 dias; contendo as modificações sugeridas pela banca examinadora e constante nos formulários de avaliação da banca. A banca examinadora foi composta pelos examinadores professores doutores listados abaixo. Os pareceres assinados seguem em sequência.

*Rodrigo Fadini*

RODRIGO FERREIRA FADINI  
Orientador

*Maria Katiane Souza Costa*

MARIA KATIANE SOUSA COSTA

Discente



*Universidade Federal do Oeste do Pará*  
**PROGRAMA DE PÓS GRADUAÇÃO EM BIODIVERSIDADE**

**Dr. RENATO PORTELA SALOMÃO**

Examinador Externo à Instituição

**Dra. RAQUEL LUIZA DE CARVALHO**

Examinadora Externa à Instituição

**Dr. FABRICIO BEGGIATO BACCARO, UFAM**

Examinador Externo à Instituição

**Dr. RODRIGO FERREIRA FADINI, UFOPA**

Presidente

**MARIA KATIANE SOUSA COSTA**

Discente

A Deus, a luz que guia meus dias  
A minha mãe Maria José, ao meu filho Lucas e  
ao meu esposo Renato, vocês são a razão de  
tudo. São minha força e determinação.

## AGRADECIMENTOS

É impossível chegar ao fim desse trabalho e não pensar em todas as pessoas que contribuíram para que eu chegasse até aqui. Agradeço primeiramente a Deus pois, sem Ele não somos capazes de superar os obstáculos que a vida nos impõe. Foi Deus que me fez ter a certeza de que estava no caminho certo e de que merecia tudo de bom que a vida tem a oferecer.

Agradeço à minha família por sempre me apoiar em todas as minhas decisões. Agradeço por todos os momentos de conflito em que estiveram ao meu lado, mesmo quando me faltou tempo para estar com vocês, e até mesmo pela paciência nos momentos mais estressantes. O amor e o apoio incondicional que vem se estendendo por toda a minha vida é o alicerce que sustenta meus sonhos e me permitem transformá-los em realidade. Em especial ao meu filho Lucas Eduardo, ao meu esposo Renato Vasconcelos e minha mãe, Maria José. Essa conquista é para vocês e por vocês, que sempre foram a razão pela qual eu busco novas oportunidades

Aos meus orientadores Rodrigo F. Fadini e Filipe M. França. Nessa jornada, vocês foram parte essencial para que eu chegasse até aqui. Sem o apoio, a compreensão e acima de tudo a paciência de vocês não teria conseguido superar os percalços que surgiram e olha que não foram poucos e vocês não desistiram de mim, minha gratidão aos dois. Agradeço as minhas amigas em especial a Vanessa Gomes, Soliane Sousa e Beatriz Souza, por estarem sempre ao meu lado, desde os momentos bons até os mais difíceis. Obrigada pelas palavras de apoio, o incentivo quando nem eu mesmo me sentia capaz de dar continuidade e concluir esse trabalho, vocês acreditaram em mim e não me deixaram desistir. As meninas do laboratório de Botânica e PPbio que me receberam com muito carinho e acreditaram em mim, meu muito obrigada. Assim como os alunos, professores e técnicos do laboratório de sementes Florestais.

Não poderia esquecer o Alisson Bezerra, Dian Carlos, Carlos Brocardo que foram meus companheiros em campo, mas que companheiros, foram amigos e me ajudaram a realizar todas as coletas. Obrigada pelo apoio e compreensão. Agradeço a UFOPA e ao PPGBEES por todo conhecimento adquirido, pelo corpo de docente que em muito contribuíram para minha formação. E por toda a infraestrutura que que possibilitou desenvolver esse trabalho. Agradeço a todos que de alguma forma me incentivaram e me ajudaram durante toda minha vida. Por fim, agradeço a mim mesma por ter sido persistente e acreditado em mim e nas palavras de incentivo de todos os que me apoiaram. Vocês tinham razão eu sou capaz. Muito obrigada.

“Existem muitas hipóteses em ciência que estão erradas. Isso é perfeitamente aceitável, eles são a abertura para achar as que estão certas”. (Carl Sagan)

“A imaginação é mais importante que a ciência, porque a ciência é limitada, ao passo que a imaginação abrange o mundo inteiro”. (Albert Einstein)

## RESUMO

Variáveis ambientais são importantes estruturantes das assembleias de besouros escarabeíneos na escala local em florestas preservadas no Neotrópico, mas qual a importância relativa dessas variáveis num contexto de mudança de uso da terra com o plantio de espécies florestais exóticas (i.e., eucalipto) adjacentes às florestas nativas não está claro. Neste estudo, perguntamos: [i] Como as assembleias de besouros serão influenciadas pela textura do solo, biomassa da serrapilheira, abertura do dossel e distância da silvicultura de eucalipto? E [ii] quais são as variáveis mais importantes para estruturar a composição de espécies de besouros? Para avaliar essas questões, utilizamos GLMs, regressões múltiplas multivariadas e análises de partição de variância hierárquica. A riqueza rarefeita de espécies de besouros diminuiu com o aumento da abertura do dossel e da porcentagem de areia do solo, mas apenas a distância da plantação de eucalipto, e não a textura do solo e a abertura do dossel, afetou significativamente a abundância de indivíduos de besouros. A compreensão dos resultados desse estudo é importante num contexto de avanço do cultivo de espécies arbóreas exóticas, que vem se espalhando por várias regiões amazônicas nas últimas décadas, influenciando de uma maneira até então desconhecida a fauna do interior das florestas preservadas adjacentes.

**Palavras-Chave:** Brasil. Cobertura do dossel. Efeito de borda. Floresta de espécies exóticas. Textura do solo.

## **ABSTRACT**

Environmental variables are important for structuring dung-beetle assemblages in mature native forests the Neotropics at the local scale. However, their relative importance in the context of land-use changes with crops of exotic forest species (i.e., eucalyptus) close to native forests is still not clear. In this study we argue: [i] how are dung-beetles assemblages influenced by environmental variables (i.e., soil texture, litter biomass, canopy opening), and distance to eucalyptus plantations? and [ii] which are the most important variables for structuring dung-beetles species composition? In order to assess such issues, we used GLMs, multivariate multiple regression and hierarchical partitioning analysis. The rarefied species richness of beetle has decreased with the increase of canopy opening and the percentage of sand in the soil. However, the distance to eucalyptus plantation alone and not soil texture or canopy opening, has significantly affected beetle abundance. Understanding the results of this study is important in a context of expansion of exotic tree plantations, which has been spreading through several Amazonian regions in the last decades, thus influencing the native fauna within native preserved forests nearby.

**Keywords:** Brazil. Canopy cover. Edge effect. Exotic species forestry. Soil texture

## LISTA DE ILUSTRAÇÕES

- Fig. 1.** Map of the study area: plots, deforested areas, native forests, and eucalyptus plantations. ----- 24
- Fig. 2.** Effect of proximity to Eucalyptus plantations on the species richness of dung beetles within undisturbed primary Amazonian forests, Pará state, Brazil. ----- 30
- Fig. 3.** Rarefied species richness of dung beetles in relation to canopy openness (a) and soil sand percentage (b) in undisturbed primary Amazonian forests under the influence of Eucalyptus plantations, Pará state, Brazil. ----- 31
- Fig. 4.** Effect of distance to Eucalyptus on the number of dung beetles individuals collected in undisturbed primary Amazonian forests, Pará state, Brazil. ----- 32
- Fig. 5.** Composition of dung beetles species due to the percentage of sand in the soil, collected in undisturbed primary Amazonian forests, Pará state, Brazil. ----- 34
- Fig. 6.** Composition of dung-beetle species due to the distance of eucalyptus plantations, collected in undisturbed primary Amazonian forests, Pará state, Brazil. ----- 35
- Fig. 7.** Composition of dung beetles species (presence-absence data) due to the percentage of sand in the soil, collected in undisturbed primary Amazonian forests, Pará state, Brazil. ---- 36
- Fig. 8.** Distribution of percentage of independent effects of different predictors of dung beetles assemblages in undisturbed primary Amazonian forests. The x-axis represents the percentage of independent effects (I%) divided by the total variance explained by the model (R<sup>2</sup>de). LL = litter weight (g), CO = canopy opening (%), ST = soil texture (% of sand) e ED = distance to eucalyptus (m). Black sticks represent significant effects ( $\alpha = 0.05$ ) established by randomized tests. Z values for the distribution produced by the 1,000 randomization of independent effects were calculated as indicators of the significance based on the 0.95 ( $Z \geq 1.65$ ) confidence interval. ----- 37

## **LISTA DE TABELAS**

**Table 1.** Species and number of individuals (N) collected in pitfalls in undisturbed forest area under the influence of eucalyptus plantations in the Jari region, Pará state, Brazilian Amazon.

----- **29**

**Table 2.** Results of multivariate multiple regression on the effects of environmental and landscape variables on the structure (relative abundance) and composition of dung beetles species (presence-absence). The values outside the parenthesis indicate the regression coefficients. Significant p-values are in bold. ----- **33**

## **LISTA DE ABREVIATURAS E SIGLAS**

Am	Clima tropical úmido
GLA	Gap Light Analyzer
GLM	General Linear Models
LECIN	Laboratório de Ecologia e Conservação de Invertebrados
PCoA	Principal Coordinate Analysis
PPBio	Programa de Pesquisa em Biodiversidade
PPGBEES	Programa de Pós-graduação em Biodiversidade
UFOPA	Universidade Federal do Oeste do Pará
VIF	Variance Inflation Factor

## SUMÁRIO

<b>1. INTRODUÇÃO GERAL -----</b>	<b>14</b>
<b>2. CAPÍTULO ÚNICO -----</b>	<b>17</b>
<b>2.1. Introduction -----</b>	<b>20</b>
<b>2.2. Materials And Methods -----</b>	<b>22</b>
2.2.1 Area of study -----	22
2.2.2. Dung-beetles sampling -----	24
2.2.3. Data collection on environmental and landscape variables -----	24
2.2.4. Statistical analysis -----	25
<b>2.3. Results -----</b>	<b>27</b>
2.3.1. Species richness, abundance, and biomass -----	29
2.3.2. Species structure and composition -----	31
2.3.3. Relative importance of explanatory variables for dung-beetles assemblages -----	35
<b>2.4. Discussion -----</b>	<b>36</b>
<b>2.5. References -----</b>	<b>40</b>

## **1. INTRODUÇÃO GERAL**

### **Proximidade com plantações de eucalipto afeta os besouros rola-bosta em florestas nativas primárias da Amazônia<sup>1</sup>**

#### **Qual o problema da pesquisa?**

A subfamília Scarabaeinae é composta por besouros coprófagos mais conhecidos como rola-bosta, que apresentam ampla diversidade e distribuição geográfica. Desempenham um papel crucial em vários processos ecológicos contribuindo para a remoção de resíduos orgânicos e ciclagem de nutrientes, fertilização e aeração do solo, dispersão secundária de sementes e controle de parasitas de vertebrados. A maioria das espécies desse grupo apresenta forte associação com as variáveis ambientais que caracterizam seus habitats (estrutura do solo, abertura da copa das árvores, temperatura e umidade relativa), sendo altamente sensíveis às variações ocorrentes no ambiente

Os besouros rola-bosta vêm sendo amplamente utilizados em pesquisas ecológicas como um grupo bioindicador de mudanças ambientais, como nos processos de fragmentação florestal, de impactos causados pela mudança do uso da terra, além de fenômenos climáticos extremos e incêndios florestais. As mudanças no uso da terra e a degradação dos habitats naturais através do desmatamento, da fragmentação e da superexploração florestal são considerados as maiores ameaças à biodiversidade global. Um exemplo desse tipo de fragmentação é a substituição de imensas áreas de floresta nativa por monoculturas de eucaliptos. Essas atividades humanas promovem a substituição de ambientes heterogêneos por habitats mais simplificados e afetam negativamente a biodiversidade e suas funções ecológicas

A maioria dos estudos que avaliam a composição de espécies dos besouros rola-bosta identificam as características do habitat e da paisagem como muito importantes para determinar quais espécies, onde e quantos besouros serão encontrados. Outra questão é que em sua maioria são pesquisas comparativas entre tipos de vegetação muito diferentes (como pastos VS. florestas preservadas). Poucos estudos, no entanto, exploram como os besouros rola-bosta são influenciados colocar por mudanças sutis dentro do mesmo ambiente e da paisagem numa escala mais local, principalmente em áreas com gradientes ambientais pouco aparentes e paisagens dominadas por florestas plantadas com eucaliptos. Como será que essas florestas de eucalipto afetam os besouros que estão localizados nas florestas preservadas adjacentes?

#### **Como a pesquisa foi realizada?**

A pesquisa foi realizada na Jarí Florestal, localizada entre os estados do Pará e Amapá. Ela surgiu da necessidade de se conhecer a influência de diferentes características do habitat como a estrutura dosolo, abertura de dossel, profundidade da serrapilheira e da proximidade com a monocultura de eucalipto sobre o número de espécies, a quantidade de indivíduos, a biomassa (peso em gramas) e composição e estrutura das comunidades de besouros rola-bosta em florestas primárias não perturbadas da Amazônia (Figura 1). Nossas perguntas foram as seguintes: [1] Como as assembleias de besouros são influenciadas pelas características do habitat (estrutura dosolo, abertura de dossel, biomassa da serrapilheira) e pela proximidade com monocultura de eucalipto? E [2] quais são as características mais importantes para determinar mudanças na composição de espécies de besouros?



Figura 1. Espécies de besouros escarabeíneos encontrados na área de estudo.  
Fotos: Filipe Machado França.

Coletamos informações dos besouros e das características do habitat em 34 locais, distribuídos dentro de uma área de floresta nativa primária circundada por plantios de eucaliptos. Em cada local avaliamos o solo, a serrapilheira, a abertura da copa das árvores e a distância até a plantação de eucalipto. Também instalamos armadilhas de queda do tipo *pittfall*, em que colocamos um porta-isca contendo fezes para atração e coleta dos besouros. A partir das informações coletadas em campo obtivemos um banco de dados que nos permitiu fazer inúmeras análises para identificar quais das características do habitat e da paisagem influenciam os besouros rola-bosta.

## **Qual a importância da pesquisa?**

Os resultados deste estudo mostram que não só características do habitat em florestas primárias influenciam na distribuição dos besouros rola-bosta, mas também características da paisagem, em específico, a distância até o plantio de eucalipto. Plantios com espécies florestais exóticas vêm se tornando cada vez mais comuns no Brasil e na Amazônia em particular, e seus efeitos sobre a fauna de florestas nativas precisam ser levados em consideração no planejamento para uso o adequado da terra.

## **Autores**

Maria Katiane S. Costa<sup>1</sup>, Rodrigo F. Fadini<sup>1</sup>, Filipe M. França<sup>2</sup>, Carlos R. Brocardo<sup>1</sup>

## **Instituição**

<sup>1</sup> Universidade Federal do Oeste do Pará, Programa de Pós-graduação em Biodiversidade, Laboratório de Ecologia e Conservação, Campus Tapajós, Rua Vera Paz, s/n, Salé, CEP 68040-255, Santarém, PA, Brasil

<sup>2</sup> University of Bristol, 24 Tyndall Avenue, Bristol, BS8 1TQ, Reino Unido.

<sup>1</sup>Autor para correspondência: E-mail: katianecosta30@gmail.com

## **Sugestões de leitura:**

Cajaiba, R. L., & da Silva, W. B. (2015). Abundância e Diversidade de Coleópteros de Solo em Fragmentos de Capoeira ao Entorno da Zona Urbana do Município de Uruará-PA, Brasil. *EntomoBrasilis*, 8(1), 30-37.

Silva, P. G. D., Vaz-de-Mello, F. Z., & Mare, R. A. D. (2011). Guia de identificação das espécies de Scarabaeinae (Coleoptera: Scarabaeidae) do município de Santa Maria, Rio Grande do Sul, Brasil. *Biota Neotropica*, 11, 329-345.

Silva, R. J., Ribeiro, H. V., Souza, M. F., & Vaz-de-Mello, F. Z. (2015). Influência da granulometria do solo na estrutura de guildas funcionais de besouros rola-bostas (Coleoptera: Scarabaeidae: Scarabaeinae) em florestas semideciduais no estado do Mato Grosso, Brasil. *Bioscience Journal*, 31(2).

## **2. CAPÍTULO ÚNICO**

Maria Katiane Sousa Costa, Filipe Machado França, Carlos Rodrigo Brocardo, Rodrigo Ferreira Fadini. (2022). Environmental variables and proximity to exotic tree plantations affect dung-beetles in forests with sandy soils in the Amazon. Manuscrito Submetido ao periódico Forest Ecology and Management. As normas indicadas para a redação de artigos pela revista estão disponíveis no link: <https://www.elsevier.com/journals/forest-ecology-and-management/0378-1127/guide-for-authors>

**Environmental variables and proximity to exotic tree plantations drive dung beetle  
communities within Amazonian undisturbed forests**

Maria Katiane Sousa Costa<sup>a</sup>, Filipe Machado França<sup>b,\*†</sup>, Carlos Rodrigo Brocardo<sup>a</sup>, Rodrigo Ferreira Fadini<sup>a,\*†</sup>

<sup>a</sup> *Universidade Federal do Oeste do Pará, Programa de Pós-graduação em Biodiversidade,  
Rua Vera Paz, S/N, 68040-255, Santarém, PA, Brazil*

<sup>b</sup> *University of Bristol, School of Biological Sciences, 24 Tyndall Avenue, Bristol, BS8  
1TQ, UK*

\*These authors contributed equally to this work.

†Corresponding authors. Email: [\(F.M.F\)](mailto:filipe.machadofranca@bristol.ac.uk) and [\(R.F.F\)](mailto:rodrigo.fadini@ufopa.edu.br)

1 ABSTRACT

2 Environmental variables are important for structuring dung-beetle assemblages in Neotropical  
3 forests. However, their relative importance in the context of land-use changes with crops of exotic  
4 forest species (i.e., *Eucalyptus* spp) close to native forests is still not clear. In this study we  
5 investigate [i] how dung beetle assemblages (i.e., rarefied species richness, biomass, abundance  
6 and species composition) respond to local environmental variables (i.e., soil texture, litter  
7 biomass, canopy opening) and proximity to eucalyptus plantations, and [ii] which metrics are the  
8 most important for structuring dung beetle assemblages We found that dung beetle species  
9 richness and abundance were higher in forests closer to *Eucalyptus* plantations, while rarefied  
10 species richness decreased with the increase of canopy opening and the percentage of sand in  
11 the soil. Our findings demonstrate the impacts of exotic tree plantations on the biodiversity of  
12 nearby native undisturbed forests, and are relevant when considering continuous expansion of  
13 exotic forest monocultures within tropical regions.

14

15 *Keywords:* Tropical forests, insect biodiversity, Edge effects, Exotic plantations, dung beetles.

## 2.1 Introduction

Habitat degradation and land-use changes are among the largest threats to global biodiversity (Dirzo and Raven, 2003; Morris, 2010). In the tropics, these disturbances are pushed mainly by the progress of monocultures and pastures (França et al. 2021) and by timber and non-timber resources exploitation within remaining forests (Barlow et al., 2016; Ceballos et al., 2015; Lewis and Maslin, 2015; Tabarelli et al., 2010). These human activities replace heterogeneous ecosystems with more simplified habitats, and negatively affect biodiversity and associated ecological functions (Braga et al., 2013; Correa et al., 2020, 2019; Martello et al., 2016).

Cultivating forest species for timber, cellulose, and coal production is a reality in several tropical countries. In the Amazon, for example, huge silviculture concessions started with rubber trees (*Hevea brasiliensis*), which replaced large areas of native forests in the early twentieth century (D'Agostini et al., 2013). As from the 1960's, two other monocultures expanded across the Amazon to meet the global timber and cellulose demands: eucalyptus (*Eucalyptus urugrandis*) and pine (*Pinus caribaea* var. *hondurensis*). The largest exotic plantations in the Brazilian Amazon are located in the states of Pará (~139,720 ha of eucalyptus), Amapá (62,880 ha of eucalyptus and 810 ha of pine), Mato Grosso (61,530 ha of eucalyptus and 10 ha of pine) and Tocantins (44,310 ha of eucalyptus and 850 ha of pine) (Pereira et al., 2010). However, the sustainable development of such practices is limited by the lack of knowledge about the impacts of these initiatives on tropical plant and animal biodiversity (Correa et al., 2016; Martello et al., 2018, 2016; Tavares et al., 2019).

The impact of silvicultural monocultures on biodiversity depends on multiple factors. For instance, planted forests with a structure similar to native forest environments can house biological communities similar to those found in undisturbed forests (Beiroz et al., 2017; Correa et al., 2020; Korasaki et al., 2013). However, although some species may benefit from landscapes with planted forests, others specialist species become extremely rare or locally

extinct (Correa *et al.*, 2016; Martello *et al.*, 2018, 2016; Tavares *et al.*, 2019; Vasconcelos *et al.*, 2009). Bremer and Farley (2010), for example, have demonstrated that biodiversity in a *Eucalyptus* monoculture was influenced by the native forest proximity, local conditions, and landscape context. Nevertheless, little is known about the influence of exotic forest monocultures on the biodiversity of neighboring native forests, which could spread the land modification effect caused by exotic tree plantations beyond its boundaries.

Dung beetles (Coleoptera: Scarabaeinae) are largely used in ecological research due to their high sensitivity to environmental changes (Feer and Hingrat, 2005; Lopes *et al.*, 2011; Silva *et al.*, 2014; França *et al.* 2016). Most studies that assess the composition of dung beetle assemblages identify environmental and landscape variables as the most important drivers (Cottenie, 2005) for both most common and rare species (Silva and Hernández, 2016). For example, areas that preserve the largest forest coverage are normally richer in dung beetle species than areas with lower canopy cover (Giménez Gómez *et al.*, 2018; Horgan, 2005).

Dung beetle diversity and biomass are markedly affected by environmental conditions of each ecosystem. Previous studies have demonstrated that the variation of clay concentration in the soil (Silva *et al.*, 2014), amount of litter (Silva and Hernández, 2016), mammal defaunation (Cullot *et al.*, 2013) and *Eucalyptus* monoculture planting (Barlow *et al.*, 2007; Gardner *et al.*, 2008) are important drivers of beetle diversity. For instance, tunneler dung beetles – i.e., species that bury their food resources immediately below or nearby the food resource – may have a disadvantage in clay soils (Griffiths *et al.*, 2015; Salomão and Iannuzzi, 2015; Silva *et al.*, 2015). Given their high sensitivity, dung beetles are widely used as indicators of environmental changes caused by forest fragmentation (Bitencourt and da Silva, 2016), land-use changes (França *et al.*, 2018; Scheffler, 2005) and extreme climate events such as droughts and forest fires (França *et al.*, 2020; Silveira *et al.*, 2016). Besides, several studies indicate that dung beetles may respond to environmental variations similarly to other biological groups (Barlow *et al.*, 2016; Edwards *et al.*, 2011). Few studies, however,

explore the way dung beetle communities are influenced by multiple local-scale microclimate conditions, particularly in Amazonian forests. Another issue is that most studies compare dung beetle communities between native and planted forests (Beiroz *et al.*, 2017; Gardner *et al.*, 2008; Silveira *et al.*, 2010). Nonetheless, to our knowledge no previous work assessed how distance to *Eucalyptus* forests affect insect biodiversity within native undisturbed tropical forests.


To fill these knowledge gaps, our study assessed the influence of environmental variables (e.g. litter depth, soil structure, and canopy opening) and proximity to *Eucalyptus* monoculture on dung beetle species richness, abundance, biomass, and species composition within undisturbed primary forests in the Brazilian Amazon. Our research aims to answer the following questions: [i] how dung beetle assemblages respond to soil texture, litter biomass, canopy opening, and distance to eucalyptus plantations? and [ii] which predictors are the most important for structuring dung beetle assemblages? Our hypotheses are that [i] dung beetle abundance, species richness and biomass will increase in forests with a larger percentage of sand in the soil, lower canopy opening, and more distant to *Eucalyptus* plantations; and that [ii] the sand soil percentage will be the most important driver of dung beetle species composition.

## 2.2 Materials and methods

### 2.2.1 Study region

Our study was carried out at Jari Florestal, a large-scale tree plantation and timber forestry concession located in Almeirim, Pará state, in the northern portion of the Brazilian Amazon ( $0^{\circ} 27'S$   $51^{\circ} 40'W$ ) (Fig. 1). Jari Florestal occupies an area of 1,259,958 ha, of which ~129,200 ha (9% of the total area) are *Eucalyptus* plantations while 545,000 ha are native forest planned to be or already selectively logged. According to Köppen's classification, the

region is under a tropical monsoon climate (Am) with a brief dry season (Andrade *et al.*, 2017). The average mean temperature is 26.4°C, while annual precipitation is between 1998.2 and 2347.7 mm. Rainfall distribution is irregular throughout the year, with more rain between March and May and less rain between September and November (Sobrinho *et al.*, 2012). Most soils in the region are rich in clay (Batista and Woessner, 1983), but sandy soils are also present (Fearnside and Rankin, 1979). The region is a mosaic of several vegetation types, especially broad-leaf equatorial forests, dense forests, sub-montane forests, high plateaus and high terraces (*terra firme* forests), open broadleaf forests, savannas, alluvial fields and woodlands (Andrade *et al.*, 2017). Primary forests within the forestry concession are subject to extractivism of non-timber forest products and low levels of subsistence hunting (Barlow *et al.*, 2010; Parry *et al.*, 2009).



**Fig. 1.** Map of the study area: plots, deforested areas, native forests, and eucalyptus plantations.

### *2.2.2 Dung beetles sampling*

Dung beetles were surveyed at 34 plots located within native undisturbed forests (Fig. 1), between June and July 2012. Each plot received a grid of six sampling points located 100 m distant from each other and 75 m away from the plot edge. Each sampling point had a pitfall trap, which consists of a 1-L plastic recipient buried at a ground level and containing 250 ml of saline solution (water, liquid detergent, and salt). A bait container with ~40g of human and pig dung (1:4 ratio, Marsh et al., 2013) was suspended above each pitfall to attract beetle. This is considered the most appropriate method to collect dung beetles (Favila and Halffter, 1997). A plastic cover protected both the pitfall and the bait container from rain. Beetles were collected after a 24-hour trap exposure in the field. All specimens were taken to the lab, where we sorted out and identified all dung beetles up to the lowest taxonomic level. To calculate the average biomass of each species, we obtained the dry weight of 15 individuals (when feasible) using a Shimatzu AY220 0.0001-g precision scale. The total community biomass was obtained by multiplying the average biomass of each species by its abundance. Collected material is deposited at the Scarabaeinae collection of Laboratório de Ecologia e Conservação de Invertebrados (LECIN) at the Federal University of Lavras, Brazil.

### *2.2.3 Data collection on environmental and landscape variables*

To assess the influence of environmental variables on dung beetles, we measured the canopy opening, litter biomass, and soil texture in the same places in which pitfall traps were installed. To quantify the canopy opening, hemispheric photographs were taken with a Nikon FC-E8 fisheye lens attached to a Nikon D40 camera which was aligned at ca ~1.20 m above soil level. One picture was taken at each sampling point and six pictures were taken per plot. Aiming to standardize the harvesting and minimize mistakes, pictures were taken early in the morning and late in the afternoon or under a cloudy sky (Zhang et al., 2005). To estimate the canopy opening based on the pictures, the software Gap Light Analyzer (GLA version 2.0, Frazer et al., 1999) was used to calculate the ratio between the total amount of open space

and the total area of the hemispherical photography (Frazer *et al.*, 1999). This methodology is widely used to assess the canopy opening in tropical forests (Gries *et al.*, 2012; Medjibe *et al.*, 2014; Niemczyk *et al.*, 2010; Silveira *et al.*, 2010). Leaf litter was collected using a 25 x 25 cm square, placed randomly about 1m from each pitfall. We gathered one litter sample for each sampling point (six samples per plot). Samples were placed to dry in a 60°C greenhouse for 96 hours and, after that, dry weighed at a 0,001 g precision Shimatzu AY220 (Shimadzu Corporation, Kyoto, Japan) scale. For the analysis, we used the average values of canopy openness and leaf litter biomass across the six sampling points per plot.

A soil sample (up to 10 cm deep) was collected in each sampling point. We mixed the six samples from each plot to represent the plot-level soil texture (clay, silt, and fractions of coarse sand). The amount of sand in the soil was considered as our measure of soil texture, as in previous studies (França *et al.*, 2018; Gries *et al.*, 2012; Griffiths *et al.*, 2015). Granulometric analysis was conducted in the Jari Celulose S.A. soil lab. The distance to *Eucalyptus* plantations was estimated by using a GPS to georeference each sampling points, while information on planting areas was obtained through a shapefile provided by Jari Florestal. The smallest distance to sampling points as to the eucalyptus planting areas was measured using software Quantum Gis (version 3.4.15) and plugin NNJoin (<https://plugins.qgis.org/plugins/NNJoin/>) which joins two vector layers (the entrance layer and the merging layer) based on closer neighbor relations and closest distance up to the *Eucalyptus* planting edge.

#### 2.2.4 Statistical analysis

Multicollinearity between predictor variables (canopy opening, litter biomass, percentage of sand and eucalyptus distance) was assessed with the variance inflation factor (VIF), using a *vif* function in *car* package (Fox and Weisberg, 2019), in software R (R Core Team, 2021). VIF value was smaller than 2.53 in all cases, indicating low multicollinearity. We, therefore, used the complete set of explanatory variables in all models (canopy opening, litter

biomass, percentage of sand, and distance to eucalyptus). Spatial autocorrelation was tested separately for the species composition and biomass data using Mantel tests (Legendre and Legendre, 1998), with 1000 permutations using the *mantel* function in the *vegan* package (Oksanen *et al.*, 2020).

To control the number of individuals' effects over the species richness observed, we also estimated the richness expected (rarefied richness) using *rarefy* function of the *vegan* package (Oksanen *et al.*, 2020). Rarefaction is an approach used to establish the standardization of sampling efforts and thus, allow a straightforward comparison of results between groups showing differences in the abundance patterns (Barlow *et al.*, 2007). Linear models with Gaussian distribution were used for continuous dependent variables (rarefied richness and biomass). For species richness and abundance variables, we used General Linear Models (GLM) with Poisson distribution and the “overdispersion” was corrected using Quasipoisson adjustment whenever necessary.

The effect of environmental variables on the composition of the dung beetles assemblage was also assessed. First, spatial autocorrelation was tested in our composition data using Mantel tests, and then, the assemblage dimension was reduced through a Principal Coordinate Analysis (PCoA) using a Bray-Curtis and Jaccard distance for abundance and presence-absence data, respectively. Species with only one or two records were removed from the analysis. Then, two axes (PCoA1) were used in a multivariate multiple regression mode, using all explanatory variables listed above. The result was plotted using the Poncho function (Dambros, 2020), which classifies the species along the gradients identified in the previous analysis.

To assess the most important drivers of dung beetle assemblages, we used a hierarchical partition analysis (Chevan and Sutherland, 1991) to compare the relative and independent importance of our four predictor variables for rarefied richness, abundance, and total biomass of the dung beetle communities. Hierarchical partitioning is a multiple regression

technique in which all possible linear models are considered together to establish the most probable predictors, while minimizing the influence of multicollinearity and providing an independent contribution from each explanatory variable. Models were assessed based on the  $R^2$  adjustment, which allowed us to interpret the independence of effects as a proportion of the variance explained by the total model. The significance of independent effects for each explanatory variable was calculated by randomized tests with 1,000 permutations (Mac Nally et al., 2002; Mac Nally and Walsh, 2004).

## 2.3 Results

We collected 3,722 dung beetles distributed in 13 genera and 49 species (Table 1). The nine most abundant species corresponded to ~85% of the individuals collected. *Oxysternon festivum* was the species with the largest number of individuals (502), followed by *Eurysternus caribaeus* (489) and *Onthophagus haematopus* (422). On the other hand, *Sulcophanaeus faunus*, *Oxysternon durantoni*, *Dichotomius worontzowi*, *Dichotomius latilobatus*, *Dichotomius* sp. 1, and *Ateuchus* sp. B presented only one individual each. A weak spatial autocorrelation was identified for dung beetle species composition and biomass ( $r = 0.46$  and  $0.24$ , respectively;  $p = 0.001$ ).


**Table 1.** Species and number of individuals (N) collected in pitfalls in undisturbed forest area under the influence of eucalyptus plantations in the Jari region, Pará state, Brazilian Amazon.

Species	N	Species	N
<i>Oxysternon festivum</i>	502	<i>Eurysternus hamaticollis</i>	9
<i>Eurysternus caribaeus</i>	489	<i>Canthidium sp. K</i>	8
<i>Onthophagus haematopus</i>	422	<i>Onthophagus bidentatus</i>	8
<i>Eurysternus caribaeus</i>	356	<i>Eurysternus balachowskyi</i>	7
<i>Dichotomius lucasi</i>	348	<i>Ateuchus sp. D</i>	6
<i>Deltochilum aff. submetallicum</i>	327	<i>Ateuchus sp. E</i>	6
<i>Ateuchus aff. murray</i>	284	<i>Canthidium aff. lenthum</i>	5
<i>Canthon triangularis</i>	278	<i>Deltochilum orbiculare</i>	5
<i>Deltochilum septemstriatum</i>	136	<i>Dichotomius subuaneus</i>	5
<i>Canthidium sp. B</i>	69	<i>Onthophagus onthochromus</i>	5
<i>Dichotomius boreus</i>	56	<i>Trichillum pauliani</i>	5
<i>Canthidium aff. deyrollei</i>	50	<i>Dichotomius imitator</i>	4
<i>Ontherus carinifrons</i>	46	<i>Canthidium sp. G</i>	3
<i>Ateuchus pauki</i>	38	<i>Canthon quadrigutatus</i>	3
<i>Deltochilum sp. B</i>	37	<i>Canthidium sp. I</i>	2
<i>Eurysternus foedus</i>	34	<i>Coprophanaeus dardanus</i>	2
<i>Uroxys sp. C</i>	30	<i>Coprophanaeus jasius</i>	2
<i>Eurysternus ventricosus</i>	27	<i>Eurysternus howdeni</i>	2
<i>Canthidium sp. D</i>	21	<i>Ateuchus sp. B</i>	1
<i>Coprophanaeus lancifer</i>	19	<i>Dichotomius latilobatus</i>	1
<i>Ateuchus sp. A</i>	15	<i>Dichotomius sp. 1</i>	1
<i>Deltochilum icarus</i>	14	<i>Dichotomius worontzowi</i>	1
<i>Ateuchus aff. connexus</i>	13	<i>Oxysternon durantoni</i>	1
<i>Ateuchus sp. F</i>	9	<i>Sulcophanaeus faunus</i>	1
<i>Canthidium sp. M</i>	9		


Litter biomass varied from 18.1 to 50.7 g ( $30.4 \pm 9.11$  g), canopy opening varied from 8.5 to 15.42% ( $11.5 \pm 1.72\%$ ), soil sand percentage varied from 62.6 to 95.2% ( $79.9 \pm 10.98\%$ ), and the closest distance up to the edge of eucalyptus plantations varied from 178 to 4,226 m ( $1,739 \pm 1,315$  m).

### 2.3.1 Species richness, abundance, and biomass

Dung beetle species richness varied from 9 to 26 ( $M_e = 16.6 \pm DP = 4.5$ ) per plot. At each 1,000 m of distance to *Eucalyptus* plantation, species richness reduced by 10% (coef = -0.0001,  $t = -2.32$ ,  $p = 0.027$ , Poisson GLM; Fig. 2). Rarefied species richness varied from 8.1 to 16 ( $M_e = 11.3 \pm DP = 1.74$ ) per plot and was negatively influenced by canopy opening ( $\beta = -0.38$ ,  $t = -2.26$ ,  $p = 0.03$ ,  $r^2 = 0.15$ ; Fig. 3a) and by soil sand percentage ( $\beta = -0.09$ ,  $t = -2.54$ ,  $p = 0.01$ ,  $r^2 = 0.18$ ; Fig. 3b).




**Fig. 2.** Effect of proximity to *Eucalyptus* plantations on the species richness of dung beetles within undisturbed primary Amazonian forests, Pará state, Brazil.



**Fig. 3.** Rarefied species richness of dung beetles in relation to canopy openness (a) and soil sand percentage (b) in undisturbed primary Amazonian forests under the influence of *Eucalyptus* plantations, Pará state, Brazil.

Dung beetle abundance varied from 38 to 356 individuals ( $M_e = 109.5 \pm DP = 63.5$ ) per plot. At each 1,000 m distant from the *Eucalyptus* plantation, abundance declined by 28% ( $\text{coef} = -0.0002$ ,  $t = -3.1$ ,  $p = 0.004$ , Poisson GLM; Fig. 4). Dung beetle biomass varied from 2.15 to 11.45 grams ( $M_e = 5.79 \pm DP = 2.6$  grams) per plot, but did not respond significantly to any explanatory variables ( $p > 0.05$ , data not shown).




**Fig. 4.** Effect of distance to *Eucalyptus* on the number of dung beetles individuals collected in undisturbed primary Amazonian forests, Pará state, Brazil.

### 2.3.2 Species structure and composition


The first two axes of PcoA explained 32% of the variation in the dung beetle assemblage structure (Table 2). The percentage of sand in the soil (Fig. 5) and the distance to *Eucalyptus* (Fig. 6) significantly influenced the assemblage structure, both to the first and to the second axis of PcoA (results confirmed by MANOVA; Table 2). *Ontherus carinifrons*, *Canthidium aff. deyrolley*, and *Canthidium sp. G* were the most predominant in sandier soils, while *Onthophagus bidentatus*, *Canthon triangularis* and *Ateuchus sp. D* occurred mainly in less sandy soils. *Ateucus sp. B* and *Trichillum pauliani* occurred mainly within a longer distance to the eucalyptus plantation, while *Canthidium aff. deyrolley*, *Deltochilum sp. B* were observed only in areas close to eucalyptus plantations. The first two axes of PcoA explain 30.4% of the variation in the composition of species, which was significantly influenced by the percentage of sand in the soil, by the percentage of canopy opening, and by the distance to *Eucalyptus* (Table 2; Fig. 7).

**Table 2.** Results of multivariate multiple regression on the effects of environmental and landscape variables on the structure (relative abundance) and composition of dung beetles species (presence-absence). The values outside the parenthesis indicate the regression coefficients. Significant p-values are in bold.

	Variance explained	% of sand	Litter	Canopy cover (%)	Distance Eucalyptus to	R <sup>2</sup> -adj
<b><u>Bray-Curtis</u></b>						
PCoA 1	17	0.009 ( <b>0.02</b> )	-0.001 (0.75)	0.008 (0.68)	0.000 ( <b>0.002</b> )	0.27
PCoA 2	15	-0.01 ( <b>0.000</b> )	-0.001 (0.55)	0.003 (0.74)	0.000 ( <b>0.01</b> )	0.78
Pillai-Trace		<b>(0.000)</b>	(0.836)	(0.901)	<b>(0.003)</b>	
<b><u>Jaccard</u></b>						
PCoA 1	20.8	0.013 ( <b>0.001</b> )	-0.001 (0.72)	0.017 (0.34)	-0.000 (0.92)	0.46
PCoA 2	9.6	-0.002 (0.40)	0.008 ( <b>0.04</b> )	-0.039 ( <b>0.01</b> )	-0.000 ( <b>0.01</b> )	0.20
Pillai-Trace		<b>(0.007)</b>	(0.122)	<b>(0.038)</b>	<b>(0.047)</b>	




**Fig. 5.** Composition of dung beetles species due to the percentage of sand in the soil, collected in undisturbed primary Amazonian forests, Pará state, Brazil.



**Plots ordered by distance to Eucalyptus (m)**


**Fig. 6.** Composition of dung-beetle species due to the distance of eucalyptus plantations, collected in undisturbed primary Amazonian forests, Pará state, Brazil.



**Fig. 7.** Composition of dung beetles species (presence-absence data) due to the percentage of sand in the soil, collected in undisturbed primary Amazonian forests, Pará state, Brazil.

### 2.3.3 Relative importance of explanatory variables for dung beetles assemblages

Hierarchical partitioning and randomization tests reinforced GLM results showing the influence of environmental variables for the dung beetle assemblages (Fig. 8). Canopy opening and soil sand content have significantly influenced the rarefied species richness, while the distance to *Eucalyptus* explained significantly and in an independent way the changes in the total abundance and biomass of dung beetles assemblages.



**Fig. 8.** Distribution of percentage of independent effects of different predictors of dung beetles assemblages in undisturbed primary Amazonian forests. The x-axis represents the percentage of independent effects (IE%) divided by the total variance explained by the model ( $R^2_{\text{de}}$ ). LL = litter weight (g), CO = canopy opening (%), ST = soil texture (% of sand) e ED = distance to eucalyptus (m). Black sticks represent significant effects ( $\alpha = 0.05$ ) established by randomized tests. Z values for the distribution produced by the 1,000 randomization of independent effects were calculated as indicators of the significance based on the 0.95 ( $Z \geq 1.65$ ) confidence interval.

## 2.4 Discussion

Although previous studies have compared biodiversity between planted forests and native forests (Beiroz *et al.*, 2019; Gardner *et al.*, 2008), we do not know studies that assess how forest monoculture plantations influence insects' fauna in nearby undisturbed forests in Amazonia or any other tropical regions. While addressing this knowledge gap, our results show the influence of proximity to *Eucalyptus* monoculture, canopy opening and soil texture on dung beetles communities from Amazonian forests. Understanding our results is important in the context of the development of exotic tree species initiatives, which has been spreading through the Amazon and other tropical forested regions over the last decades (Barlow *et al.*, 2007; De Oliveira *et al.*, 2019; Aguiar *et al.*, 2014; Benami *et al.*, 2018; Campos *et al.*, 2021; Homma and Vieira, 2012; Stephens and Wagner, 2007).

The rarefied species richness of dung-beetle species declined with the increase of the canopy opening and the percentage of sand in the soil. In little less than 30% increase of sand percentage in the soil (62.6 – 95.2), on average there was a reduction of three species, about the same amounts of species lost with the increase of ~7% in the canopy opening. Dung

beetles are sensitive to alterations in soil texture (Griffiths *et al.*, 2015; Nichols *et al.*, 2007; Silva *et al.*, 2015) and to the canopy structure of the forest (Gómez-Cifuentes *et al.*, 2019; Horgan, 2005). Salomão *et al.* (2022) have shown that the increase in the percentage of clay positively affects the number of tunneler beetle species, but not of roller or resident beetles. However, they have studied much less sandy soils ( $20.8\pm10.4\%$ ) than our study did ( $80\pm11\%$ ). Few species seem to tolerate very sandy soils, which retain less moisture and have a low compaction capacity, which results, respectively, in the desiccation of food resources and makes it more difficult to build nesting tunnels (Lumaret *et al.*, 1991; Sowig, 1995).

In our area of study, only four species (*Ontherus carinifrons*, *Canthidium aff. deyrolleyi*, *Canthidium sp.*, and *Canthidium aff. lendum*) seem to tolerate extremely sandy soils, since these species were only recorded in areas with more than 70% of the amount of sand. Sandy soils may become even more desiccated in larger canopy openings, since these openings are responsible for regulating the microclimate conditions of the soil and the understory, such as temperature, dampness, and radiant heat in the environment (Davis *et al.*, 2001; Tuff *et al.*, 2016). Perhaps because of this, a subtle increase in canopy opening may have significantly and negatively affected the species richness of beetle in our study, a similar pattern to that found in several studies (da Silva, 2011; Gardner *et al.*, 2008; Marinoni and Gelho, 2006; Navarrete and Halffter, 2008; Nichols *et al.*, 2007; Qie *et al.*, 2011; Viegas *et al.*, 2014)

Despite the effects on species richness, the distance to *Eucalyptus* plantations alone, but not soil texture and canopy opening, have significantly affected the abundance of individuals. By analyzing each functional guild separately, the distance to eucalyptus plantations was the only variable that influenced tunnelers and resident beetles, but not roller ones (data not shown). Beiroz *et al.* (2017) have shown a strong effect of soil on the total abundance of dung beetles in forests in the same region. This difference, however, may be a result of the spatial scale of the soil sample in that study (ca. 27 km on average) compared to ours ( $1.7\pm1.07$  km). The relative effects of the distance to eucalyptus plantations on the

abundance of dung beetles were demonstrated both by GLM and by the analysis of the hierarchical partition of variance. This demonstrates that reforestation of exotic species may influence biodiversity in surrounding native forests, thus promoting an increase of opportunistic species. This is worrisome if we consider that the new Brazilian Forest Legislation allows that the reclamation of 50% of legal reserves total areas in rural properties be made with exotic species (Brasil, 2012). Besides the risks related to invasion of exotic plant species (Zenni and Ziller, 2011), our study shows that there was not enough time for dung beetle assemblages to restructure after the impact, even decades after the eucalyptus plantation was concluded.

When we analyze the independent effects of each variable, our results demonstrate that the distance of eucalyptus planting was the only variable related to the abundance and total biomass of beetle communities (Fig. 8). The effects of *Eucalyptus* planting or other forest monocultures on dung beetles may occur directly or indirectly. For example, the proximity of exotic forest monocultures may reduce soil moisture due to high hydric demand (Poor and Fries, 1985) and also because the canopy allows the entrance of more light and heat, thus jeopardizing nesting of species most sensible to microclimate alterations (Nichols *et al.*, 2007). It is also possible that the vertebrates' fauna is altered in the vicinities of the *Eucalyptus* plantation (Pattanavibool and Dearden, 2002; Terborgh *et al.*, 2001; Umapathy and Kumar, 2000), thus reducing the availability of resources for dung beetles. In Brazil, forested planted with *Eucalyptus* may have a large diversity of mammal species (Piña *et al.*, 2019), however, due to a uniform and scattered canopy, they cannot host large populations and diversity of large size primates (Gardner *et al.*, 2008), which is one of the groups that most produced faeces used by dung beetles as a food resource in Neotropic forests (Estrada *et al.*, 1993; Feer, 1999).

Lastly, our study corroborates previous studies demonstrating the influence of forest degradation on tropical biodiversity (Barlow *et al.*, 2016; França *et al.*, 2018). Our results have shown, in particular, the importance of conservation strategies to mitigate the edge effects that exotic forest monocultures may cause in the biodiversity of nearby native forests.

### **CrediT authorship contribution statement**

**Maria Katiane S. Costa:** Writing - original draft, Writing - review & editing. **Filipe M. França:** Supervision, Conceptualization, Methodology, Investigation, Resources, Data Curation, Formal analysis, Writing - review & editing, Project administration. **Carlos R. Brocardo:** Writing - review & editing, Formal analysis. **Rodrigo F. Fadini:** Supervision, Conceptualization, Formal analysis, Writing - review & editing.

### **Declaration of Competing Interest**

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

**Acknowledgements:** The authors are grateful to Jari Florestal for logistical support. We thank the field assistants Edivar Dias Correa and Jucelino Alves dos Santos. We are grateful to Fernando Z. Vaz-de-Mello for help in the identification of specimen material. We also thank Julio Louzada, Vanesca Korasaki, Juliana Silveira and Jos Barlow for support during sampling design and fieldwork. This research was supported by grants from the MCTI/CNPq/FAPs (Nº.34/2012), CNPq-PELD site 23 (403811/2012-0), and PROCAD-AM (Nº 88881.314420/2019-01). M.K.S.C. received a master fellowship from CAPES. F.M.F. received fellowships from CAPES (BEX 5528/13-5; DTI-B-B 383744/2015-6) and acknowledges funding provided by BNP Paribas Foundation's Climate and Biodiversity Initiative (Project Bioclimate); and the Brazilian Research Council (CNPq-CAPES; Prevfogo-IBAMA 441949/2018-5 [Sem-Flama], MCIC 420254/2018-8 [Resflora] and 441659/2016-0 [PELD-RAS]). C.R.B. received a post-doctoral fellowship from CAPES (Finance Code 001).

## 2.5. References

- Aguiar Jr, A., Barbosa, R. I., Barbosa, J. B., Mourão Jr, M. 2014. Invasion of *Acacia mangium* in Amazonian savannas following planting for forestry. *Plant Ecol. Divers.* 7, 359–369. <https://doi.org/10.1080/17550874.2013.771714>
- Andrade, V.M.S., Cordeiro, I.M.C.C., Schwarts, G., Rangel-Vasconcelos, L.G.T., Oliveira, F.A., 2017. Considerações sobre clima e aspectos edafoclimáticos da mesorregião nordeste paraense, in: Embrapa. pp. 59–96.
- Barlow, J., Gardner, T.A., Araujo, I.S., Ávila-Pires, T.C., Bonaldo, A.B., Costa, J.E., Esposito, M.C., Ferreira, L. V., Hawes, J., Hernandez, M.I.M., Hoogmoed, M.S., Leite, R.N., Lo-Man-Hung, N.F., Malcolm, J.R., Martins, M.B., Mestre, L.A.M., Miranda-Santos, R., Nunes-Gutjahr, A.L., Overal, W.L., Parry, L., Peters, S.L., Ribeiro, M.A., Da Silva, M.N.F., Da Silva Motta, C., Peres, C.A., 2007. Quantifying the biodiversity value of tropical primary, secondary, and plantation forests. *Proc. Natl. Acad. Sci. U. S. A.* 104, 18555–18560. <https://doi.org/10.1073/pnas.0703333104>
- Barlow, J., Lennox, G.D., Ferreira, J., Berenguer, E., Lees, A.C., Nally, R. Mac, Thomson, J.R., Ferraz, S.F.D.B., Louzada, J., Oliveira, V.H.F., Parry, L., Ribeiro De Castro Solar, R., Vieira, I.C.G., Aragão, L.E.O.C., Begotti, R.A., Braga, R.F., Cardoso, T.M., Jr, R.C.D.O., Souza, C.M., Moura, N.G., Nunes, S.S., Siqueira, J.V., Pardini, R., Silveira, J.M., Vaz-De-Mello, F.Z., Veiga, R.C.S., Venturieri, A., Gardner, T.A., 2016. Anthropogenic disturbance in tropical forests can double biodiversity loss from deforestation. *Nature* 535, 144–147. <https://doi.org/10.1038/nature18326>
- Barlow, J., Silveira, J.M., Cochrane, M.A., 2010. Fire scars on amazonian trees: exploring the cryptic fire history of the ilha de maracá. *Biotropica* 42, 405–409. <https://doi.org/10.1111/j.1744-7429.2010.00646.x>
- Batista, M.P., Woessner, R.A., 1983. Comparação do crescimento de quatro espécies exóticas em solos podzólicos na amazônia. *Rev. Floresta* 29–35.
- Beiroz, W., Barlow, J., Slade, E.M., Borges, C., Louzada, J., Sayer, E.J., 2019. Biodiversity in tropical plantations is influenced by surrounding native vegetation but not yield: a case study with dung beetles in Amazonia. *For. Ecol. Manage.* 444, 107–114. <https://doi.org/10.1016/j.foreco.2019.04.036>
- Beiroz, W., Slade, E.M., Barlow, J., Silveira, J.M., Louzada, J., Sayer, E., 2017. Dung beetle community dynamics in undisturbed tropical forests: implications for ecological evaluations of land-use change. *Insect Conserv. Divers.* 10, 94–106. <https://doi.org/10.1111/icad.12206>
- Benami, E., Curran, L.M., Cochrane, M., Venturieri, A., Franco, R., Kneipp, J., Swartos, A., 2018. Oil palm land conversion in Pará, Brazil, from 2006-2014: evaluating the 2010 Brazilian Sustainable Palm Oil Production Program. *Environ. Res. Lett.* 13. <https://doi.org/10.1088/1748-9326/aaa270>
- Bitencourt, B.S., da Silva, P.G., 2016. Forest regeneration affects dung beetle assemblages

(Coleoptera: Scarabaeinae) in the southern Brazilian Atlantic Forest. J. Insect Conserv. 20, 855–866. <https://doi.org/10.1007/s10841-016-9917-3>

Braga, R.F., Korasaki, V., Andresen, E., Louzada, J., 2013. Dung beetle community and functions along a habitat-disturbance gradient in the Amazon: a rapid assessment of ecological functions associated to biodiversity. PLoS One 8, e57786. <https://doi.org/10.1371/JOURNAL.PONE.0057786>

Bremer, L. L. and, & Farley, K. A. (2010). Does plantation forestry restore biodiversity or create green deserts ? A synthesis of the effects of land-use transitions on plant species richness. Biodivers Conserv, 19, 3893–3915. <https://doi.org/10.1007/s10531-010-9936-4>

Campos, M.S., Adami, M., Araújo, A.C. de, 2021. Análise do albedo de superfície da palma de óleo e diferentes usos e coberturas do solo no Leste da Amazônia. Rev. Bras. Meteorol. 36, 15–21. <https://doi.org/10.1590/0102-77863540071>

Ceballos, G., Ehrlich, P.R., Barnosky, A.D., García, A., Pringle, R.M., Palmer, T.M., 2015. Accelerated modern human-induced species losses: entering the sixth mass extinction. Sci. Adv. 1. <https://doi.org/10.1126/sciadv.1400253>

Chevan, A., Sutherland, M., 1991. Hierarchical partitioning. The American Statistician, 452, 90-96.

Correa, C.M.A., Braga, R.F., Puker, A., Korasaki, V., 2019. Patterns of taxonomic and functional diversity of dung beetles in a human-modified variegated landscape in Brazilian Cerrado. J. Insect Conserv. 23, 89–99. <https://doi.org/10.1007/s10841-018-00118-6>

Correa, C.M.A., Puker, A., Abot, A.R., 2020. Impacts of exotic pasture establishment on dung beetle assemblages (Coleoptera: Scarabaeidae: Scarabaeinae) in the brazilian cerrado. Environ. Entomol. 49, 1335–1344. <https://doi.org/10.1093/ee/nvaa132>

Correa, C.M.A., Puker, A., Ferreira, K.R., Cristaldo, C.M., Ferreira, F.N.F., Abot, A.R., Korasaki, V., 2016. Using dung beetles to evaluate the conversion effects from native to introduced pasture in the Brazilian Pantanal. J. Insect Conserv. 20, 447–456. <https://doi.org/10.1007/s10841-016-9877-7>

Cottenie, K., 2005. Integrating environmental and spatial processes in ecological community dynamics. Ecol. Lett. 8, 1175–1182. <https://doi.org/10.1111/j.1461-0248.2005.00820.x>

Culot, L., Bovy, E., Vaz-de-Mello, F. Z., Guevara, R., & Galetti, M. 2013. Selective defaunation affects dung beetle communities in continuous Atlantic rainforest. Biol. Conserv. 163, 79–89. <https://doi.org/10.1016/j.biocon.2013.04.004>

D'Agostini, S., Bacilieri, S., Vitiello, N., Hojo, H., Bilynskyj, M.C.V., Filho, A.B., Rebouças, M.M., 2013. Ciclo econômico da borracha-seringueira hevea brasiliensis (hbk) m. Arg. Páginas do Inst. Biol. 9, 6–14.

da Silva, P.G., 2011. Dung beetles (coleoptera: Scarabaeidae: Scarabaeinae) of two non-native habitats

- in Bagé, Rio Grande do Sul, Brazil. Zool. Stud. 50, 546–559.
- da Silva, P.G., Hernández, M.I.M., 2016. Spatial variation of dung beetle assemblages associated with forest structure in remnants of southern Brazilian Atlantic Forest. Rev. Bras. Entomol. 60, 73–81. <https://doi.org/10.1016/j.rbe.2015.11.001>
- Dambros, C. S. 2020. csdambros/R-functions: First release. [https://github.com/csdambros/R-functions/tree/v1.0]. DOI: 10.5281/zenodo.3784397.
- Davis, A.J., Holloway, J.D., Huijbregts, H., Krikken, J., Kirk-Spriggs, A.H., Sutton, S.L., 2001. Dung beetles as indicators of change in the forests of northern Borneo. J. Appl. Ecol. 38, 593–616. <https://doi.org/10.1046/j.1365-2664.2001.00619.x>
- De Oliveira, C.P., Francelino, M.R., Daher, M., De Araújo, E.J.G., De Souza Sanches, L., De Andrade, K.D.C., De Campos, J.S.N., 2019. Estimation of the aboveground biomass and carbon stocks in open Brazilian Savannah developed on sandy soils. Carbon Balance Manag. 14, 1–10. <https://doi.org/10.1186/s13021-019-0121-0>
- Dirzo, R., Raven, P.H., 2003. Global state of biodiversity and loss. Annu. Rev. Environ. Resour. 28, 137–167. <https://doi.org/10.1146/annurev.energy.28.050302.105532>
- Edwards, D.P., Larsen, T.H., Docherty, T.D.S., Ansell, F.A., Hsu, W.W., Derhé, M.A., Hamer, K.C., Wilcove, D.S., 2011. Degraded lands worth protecting: the biological importance of Southeast Asia's repeatedly logged forests. Proc. Royal Soc. B: Biol. Sci. 278, 82–90. <https://doi.org/10.1098/rspb.2010.1062>
- Favila, M.E., Halffter, G., 1997. The use of indicator groups for measuring biodiversity. Acta Zool. Mex. 72, 1–25.
- Fearnside, P. M., Rankin, J. M. 1979. Avaliação da Jari Florestal e Agropecuária Ltda. como modelo para o desenvolvimento da Amazônia. Acta Amaz. 9, 609–615.
- Feer, F., Hingrat, Y., 2005. Effects of forest fragmentation on a dung beetle community in French Guiana. Conserv. Biol. 19, 1103–1112. <https://doi.org/10.1111/j.1523-1739.2005.00087.x>
- Fox, J., Weisberg, S. 2019. An {R} companion to applied regression, 3<sup>rd</sup> Edition. Thousand Oaks CA: Sage. URL: <https://socialsciences.mcmaster.ca/jfox/Books/Companion/>
- França, F. M., Barlow, J., Araújo, B., Louzada, J. 2016. Does selective logging stress tropical forest invertebrates? Using fat stores to examine sublethal responses in dung beetles. Ecology and Evolution, 6:23, 8526-8533. <https://doi.org/10.1002/ece3.2488>
- França, F., Louzada, J., Barlow, J., 2018. Selective logging effects on 'brown world' faecal-detritus pathway in tropical forests: a case study from Amazonia using dung beetles. For. Ecol. Manage. 410, 136–143. <https://doi.org/10.1016/j.foreco.2017.12.027>

- França, F. M., Benkwitt, C.E., Peralta, G., Robinson, J.P.W., Graham, N.A.J., Tylianakis, J.M., Berenguer, E., Lees, A.C., Ferreira, J., Louzada, J., Barlow, J., 2020. Climatic and local stressor interactions threaten tropical forests and coral reefs. *Philos. Trans. R. Soc. B Biol. Sci.* <https://doi.org/10.1098/rstb.2019.0116>
- França, F. M., Ferreira, J., Vaz-de-Mello, F.Z., Maia, L.F., Berenguer, E., Ferraz Palmeira, A., Fadini, R., Louzada, J., Braga, R., Hugo Oliveira, V., Barlow, J., 2020. El Niño impacts on human-modified tropical forests: consequences for dung beetle diversity and associated ecological processes. *Biotropica* 52, 252–262. <https://doi.org/10.1111/btp.12756>
- França, F. M., Solar, R., Lees, A. C., Martins, L. P., Berenguer, E., Barlow, J. 2021. Reassessing the role of cattle and pasture in Brazil's deforestation: A response to "Fire, deforestation, and livestock: When the smoke clears". *Land Use Policy*, 108, 105195. <https://doi.org/10.1016/j.landusepol.2020.105195>
- Frazer, G.W., Lertzman, K.P., Sallaway, P., Marinakis, D., 1999. GLA Version 2.0, Users manual and program documentation. Version 2.0, 36.
- Gardner, T.A., Hernández, M.I.M., Barlow, J., Peres, C.A., 2008. Understanding the biodiversity consequences of habitat change: the value of secondary and plantation forests for neotropical dung beetles. *J. Appl. Ecol.* 45, 883–893. <https://doi.org/10.1111/j.1365-2664.2008.01454.x>
- Giménez Gómez, V.C., Verdú, J.R., Guerra Alonso, C.B., Zurita, G.A., 2018. Relationship between land uses and diversity of dung beetles (Coleoptera: Scarabaeinae) in the southern Atlantic forest of Argentina: which are the key factors? *Biodivers. Conserv.* 27, 3201–3213. <https://doi.org/10.1007/s10531-018-1597-8>
- Gómez-Cifuentes, A., Giménez Gómez, V.C., Moreno, C.E., Zurita, G.A., 2019. Tree retention in cattle ranching systems partially preserves dung beetle diversity and functional groups in the semideciduous Atlantic forest: the role of microclimate and soil conditions. *Basic Appl. Ecol.* 34, 64–74. <https://doi.org/10.1016/j.baae.2018.10.002>
- Gries, R., Louzada, J., Almeida, S., Macedo, R., Barlow, J., 2012. Evaluating the impacts and conservation value of exotic and native tree afforestation in Cerrado grasslands using dung beetles. *Insect Conserv. Divers.* 5, 175–185. <https://doi.org/10.1111/j.1752-4598.2011.00145.x>
- Griffiths, H. M., Louzada, J., Bardgett, R.D., Beiroz, W., França, F., Tregidgo, D., Barlow, J., 2015. Biodiversity and environmental context predict dung beetle-mediated seed dispersal in a tropical forest field experiment. *Ecology* 96, 1607–1619. <https://doi.org/10.1890/14-1211.1>
- Homma, A., Vieira, I., 2012. Colóquio Sobre Dendezeiro: Prioridades De Pesquisas Econômicas, Sociais E Ambientais Na Amazônia. *Amaz. ciência e Desenvolv.* 8, 79–90.
- Horgan, F.G., 2005. Effects of deforestation on diversity, biomass and function of dung beetles on the

eastern slopes of the Peruvian Andes. For. Ecol. Manage. 216, 117–133.  
<https://doi.org/10.1016/j.foreco.2005.05.049>

Korasaki, V., Braga, R.F., Zanetti, R., Moreira, F.M.S., Vaz-de-Mello, F.Z., Louzada, J., 2013. Conservation value of alternative land-use systems for dung beetles in Amazon: Valuing traditional farming practices. Biodivers. Conserv. 22, 1485–1499. <https://doi.org/10.1007/s10531-013-0487-3>

Legendre, P., Legendre, L. 1998. Numerical ecology. Second English edition. Elsevier, Amsterdam.

Lewis, S.L., Maslin, M.A., 2015. Defining the Anthropocene. Nature 519, 171–180.  
<https://doi.org/10.1038/nature14258>

Lopes, J., Korasaki, V., Catelli, L.L., Marçal, V.V.M., Nunes, M.P.B.P., 2011. A comparison of dung beetle assemblage structure (Coleoptera: Scarabaeidae: Scarabaeinae) between an Atlantic forest fragment and adjacent abandoned pasture in Paraná, Brazil. Zoologia 28, 72–79.  
<https://doi.org/10.1590/S1984-46702011000100011>

Lumaret, J.-P., Kirk, A.A., Hanski, I., Cambefort, Y., 1991. South temperate dung beetles. Dung Beetle Ecol. 97–115.

Mac Nally, R., Parkinson, A., Horrocks, G., Young, M., 2002. Current loads of coarse woody debris on southeastern Australian floodplains: Evaluation of change and implications for restoration. Restor. Ecol. 10, 627–635. <https://doi.org/10.1046/j.1526-100X.2002.01043.x>

Mac Nally, R., Walsh, C.J. 2004. Hierarchical partitioning public-domain software. Biodiv. Conserv. 13, 659–660.

Marinoni, R.C., Ganho, N.G., 2006. A diversidade diferencial beta de Coleoptera (Insecta) em uma paisagem antropizada do Bioma Araucária. Rev. Bras. Entomol. 50, 64–71.  
<https://doi.org/10.1590/S0085-56262006000100009>

Marsh, C.J., Louzada, J., Beiroz, W., Ewers, R.M., 2013. Optimising bait for pitfall trapping of Amazonian dung beetles (Coleoptera: Scarabaeinae). PLoS One 8, 73–147.

Martello, F., Andriolli, F., de Souza, T.B., Dodonov, P., Ribeiro, M.C., 2016. Edge and land use effects on dung beetles (Coleoptera: Scarabaeidae: Scarabaeinae) in Brazilian cerrado vegetation. J. Insect Conserv. 20, 957–970. <https://doi.org/10.1007/s10841-016-9928-0>

Martello, F., Bello, F. de, Morini, M.S. de C., Silva, R.R., Souza-Campana, D.R. de, Ribeiro, M.C., Carmona, C.P., 2018. Homogenization and impoverishment of taxonomic and functional diversity of ants in *Eucalyptus* plantations. Sci. Rep. 8. <https://doi.org/10.1038/S41598-018-20823-1>

Medjibe, V.P., Poulsen, J.R., Clark, C.J., Mbani, O.A., 2014. Natural regeneration of selected timber species in the Republic of Congo. Afr. J. Ecol. 52, 552–563. <https://doi.org/10.1111/aje.12167>

Morris, R.J., 2010. Anthropogenic impacts on tropical forest biodiversity: A network structure and

ecosystem functioning perspective. Philos. Trans. R. Soc. B Biol. Sci. <https://doi.org/10.1098/rstb.2010.0273>

Navarrete, D., Halffter, G., 2008. Dung beetle (Coleoptera: Scarabaeidae: Scarabaeinae) diversity in continuous forest, forest fragments and cattle pastures in a landscape of Chiapas, Mexico: the effects of anthropogenic changes. Biodivers. Conserv. 17, 2869–2898. <https://doi.org/10.1007/s10531-008-9402-8>

Nichols, E., Larsen, T., Spector, S., Davis, A.L., Escobar, F., Favila, M., Vulinec, K., 2007. Global dung beetle response to tropical forest modification and fragmentation: a quantitative literature review and meta-analysis. Biol. Conserv. 137, 1–19. <https://doi.org/10.1016/j.biocon.2007.01.023>

Niemczyk, T., Deppe, F., Huebl, H., Menzel, E.P., Hocke, F., Schwarz, M.J., Garcia-Ripoll, J.J., Zueco, D., Hümmer, T., Solano, E., Marx, A., Gross, R., 2010. Circuit quantum electrodynamics in the ultrastrong-coupling regime. Nat. Phys. 6, 772–776. <https://doi.org/10.1038/nphys1730>

Oksanen, J., Blanchet, F.G., Friendly, M., Kindt, R., Legendre, P., McGlinn, D., Minchin, P.R., O'hara, R.B., Simpson, G.L., Solymos, P., Henry, M., Stevens, H., Szoebs, E., Maintainer, H.W., 2020. vegan: Community Ecology Package. R package version 2.5-7. <https://CRAN.R-project.org/package=vegan>.

Parry, L., Barlow, J., Peres, C.A., 2009. Hunting for sustainability in tropical secondary forests. Conserv. Biol. 23, 1270–1280. <https://doi.org/10.1111/j.1523-1739.2009.01224.x>

Pattanavibool, A., Dearden, P., 2002. Fragmentation and wildlife in montane evergreen forests, northern Thailand. Biol. Conserv. 107, 155–164. [https://doi.org/10.1016/S0006-3207\(02\)00056-3](https://doi.org/10.1016/S0006-3207(02)00056-3)

Pereira, D., Santos, D., Vedovedo, M., Guimarães, J., Veríssimo, A., 2010. Fatos florestais da Amazônia, Imazon.

Poor, M. E, D, I, U.M., Fries, C., 1985. The ecological effects of *Eucalyptus*. Food Agric. Organ. United Nations 92.

Qie, L., Lee, T.M., Sodhi, N.S., Lim, S.L.H., 2011. Dung beetle assemblages on tropical land-bridge islands: Small island effect and vulnerable species. J. Biogeogr. 38, 792–804. <https://doi.org/10.1111/j.1365-2699.2010.02439.x>

Salomão, R.P., Iannuzzi, L., 2015. Dung beetle (Coleoptera, Scarabaeidae) assemblage of a highly fragmented landscape of Atlantic forest: from small to the largest fragments of northeastern Brazilian region. Rev. Bras. Entomol. 59, 126–131. <https://doi.org/10.1016/j.rbe.2015.03.008>

Salomão, R.P., Pires, D. de A., Baccaro, F.B., Schiatti, J., Vaz-de-Mello, F.Z., Lima, A.P., Magnusson, W.E., 2022. Water table level and soil texture are important drivers of dung beetle diversity in Amazonian lowland forests. Appl. Soil Ecol. 170. <https://doi.org/10.1016/j.apsoil.2021.104260>

Sobrinho, T.R.G., Quintairos, M.V.R., Gomes, R.C.A.S.R., Santana, E.J.M., 2012. Classificação climática conforme a metodologia Köppen do município de Laranjal do Jari/Amapá/Brasil. CONNEPI 1–7.

Scheffler, P.Y., 2005. Dung beetle (Coleoptera: Scarabaeidae) diversity and community structure across three disturbance regimes in eastern Amazonia. *J. Trop. Ecol.* 21, 9–19. <https://doi.org/10.1017/S0266467404001683>

Silva, J.S., Ribeiro, H. V., Souza, M.F., Vaz-De-Mello, F.Z., 2015. Influência da granulometria do solo na estrutura de guildas funcionais de besouros rola-bostas (Coleoptera: Scarabaeidae: Scarabaeinae) em florestas semideciduais no estado do Mato Grosso, Brasil. *Biosci.J.* 31, 601–612. <https://doi.org/10.14393/BJ-v31n1a2015-23525>

Silva, R.J., Coletti, F., Costa, D.A., Vaz-De-Mello, F.Z., 2014. Rola-bostas (Coleoptera: Scarabaeidae: Scarabaeinae) de florestas e pastagens no sudoeste da Amazônia brasileira: Levantamento de espécies e guildas alimentares. *Acta Amaz.* 44, 345–352. <https://doi.org/10.1590/1809-4392201304472>

Silveira, J.M., Barlow, J., Louzada, J., Moutinho, P., 2010. Factors affecting the abundance of leaf-litter arthropods in unburned and thrice-burned seasonally-dry amazonian forests. *PLoS One* 5, 1–7. <https://doi.org/10.1371/journal.pone.0012877>

Silveira, J.M., Louzada, J., Barlow, J., Andrade, R., Mestre, L., Solar, R., Lacau, S., Cochrane, M.A., 2016. A multi-taxa assessment of biodiversity change after single and recurrent wildfires in a Brazilian Amazon Forest. *Biotropica* 48, 170–180. <https://doi.org/10.1111/btp.12267>

Sowig, P., 1995. Habitat selection and offspring survival rate in three paracoprid dung beetles: the influence of soil type and soil moisture. *Ecography* 18, 147–154.

Stephens, S.S., Wagner, M.R., 2007. Forest plantations and biodiversity: a fresh perspective. *J. For.* 105, 307–313. <https://doi.org/10.1093/jof/105.6.307>

Tabarelli, M., Aguiar, A.V., Ribeiro, M.C., Metzger, J.P., Peres, C.A., 2010. Prospects for biodiversity conservation in the Atlantic Forest: Lessons from aging human-modified landscapes. *Biol. Conserv.* 143, 2328–2340. <https://doi.org/10.1016/j.biocon.2010.02.005>

Tavares, A., Beiroz, W., Fialho, A., Frazão, F., Macedo, R., Louzada, J., Audino, L., 2019. Eucalyptus plantations as hybrid ecosystems: implications for species conservation in the Brazilian Atlantic forest. *For. Ecol. Manage.* 433, 131–139. <https://doi.org/10.1016/j.foreco.2018.10.063>

Terborgh, J., Lopez, L., V, Percy Nuñez, Rao, M., Orihuella, G., Riveros, M., Ascanio, R., Adler, G.H., Thomas, D., Balbas, L., Terborgh, J., Lopez, L., V, Percy Nunez, Rao, M., Shahabuddin, G., Orihuella, G., Riveros, M., Ascanio, R., Adler, G.H., 2001. Ecological meltdown in predator-free forest fragments. *Science* 294, 1923–1926.

- Tuff, K.T., Tuff, T., Davies, K.F., 2016. A framework for integrating thermal biology into fragmentation research. *Ecol. Lett.* 19, 361–374. <https://doi.org/10.1111/ele.12579>
- Umapathy, G., Kumar, A., 2000. The occurrence of arboreal mammals in the rain forest fragments in the Anamalai Hills, South India. *Biol. Conserv.* 92, 311–319. [https://doi.org/10.1016/S0006-3207\(99\)00097-X](https://doi.org/10.1016/S0006-3207(99)00097-X)
- Vasconcelos, H.L., Pacheco, R., Silva, R.C., Vasconcelos, P.B., Lopes, C.T., Costa, A.N., Bruna, E.M., 2009. Dynamics of the leaf-litter arthropod fauna following fire in a neotropical woodland savanna. *PLoS One* 4, 1–9. <https://doi.org/10.1371/journal.pone.0007762>
- Viegas, G., Stenert, C., Schulz, U.H., Maltchik, L., 2014. Dung beetle communities as biological indicators of riparian forest widths in southern Brazil. *Ecol. Indic.* 36, 703–710. <https://doi.org/10.1016/j.ecolind.2013.09.036>
- Zenni, R. D., Ziller, S. R., 2011. An overview of invasive plants in Brazil. *Braz. J. Bot.* 34, 431–446
- Zhang, Y., Chen, J.M., Miller, J.R., 2005. Determining digital hemispherical photograph exposure for leaf area index estimation. *Agric. For. Meteorol.* 133, 166–181. <https://doi.org/10.1016/j.agrformet.2005.09.009>